



Oral Contraceptives and Cancer Risk

What are oral contraceptives?

Oral contraceptives (birth control pills) are hormone-containing medications that are taken by mouth to prevent pregnancy. They prevent pregnancy by inhibiting ovulation and also by preventing sperm from penetrating through the cervix.

By far the most commonly prescribed type of oral contraceptive in the United States contains synthetic versions of the natural female hormones estrogen and progesterone. This type of birth control pill is often called a combined oral contraceptive. Another type of oral contraceptive, sometimes called the mini pill, contains only progestin, which is a man-made version of progesterone.

What is known about the relationship between oral contraceptive use and cancer?

Nearly all the research on the link between oral contraceptives and cancer risk comes from observational studies, both large prospective cohort studies and population-based case-control studies. Data from observational studies cannot definitively establish that an exposure—in this case, oral contraceptives—causes (or prevents) cancer. That is because women who take oral contraceptives may differ from those who don't take them in ways other than their oral contraceptive use, and it is possible that these other differences—rather than oral contraceptive use—are what explains their different cancer risk.

Overall, however, these studies have provided consistent evidence that the risks of breast and cervical cancers are increased in women who use oral contraceptives, whereas the risks of endometrial, ovarian, and colorectal cancers are reduced (1–3).

- **Breast cancer:** An analysis of data from more than 150,000 women who participated in 54 epidemiologic studies showed that, overall, women who had ever used oral contraceptives had a slight (7%) increase in the relative risk of breast cancer compared with women who had never used oral contraceptives. Women who were currently using oral contraceptives had a 24% increase in risk that did not increase with the duration of use. Risk declined after use of oral contraceptives stopped, and no risk increase was evident by 10 years after use had stopped (4).

A 2010 analysis of data from the Nurses' Health Study, which has been following more than

116,000 female nurses who were 24 to 43 years old when they enrolled in the study in 1989, also found that participants who used oral contraceptives had a slight increase in breast cancer risk (5, 6). However, nearly all of the increased risk was seen among women who took a specific type of oral contraceptive, a "triphasic" pill, in which the dose of hormones is changed in three stages over the course of a woman's monthly cycle. An elevated risk associated with specific triphasic formulations was also reported in a nested case-control study that used electronic medical records to verify oral contraceptive use (7).

In 2017, a large prospective Danish study reported breast cancer risks associated with more recent formulations of oral contraceptives (8). Overall, women who were using or had recently stopped using oral combined hormone contraceptives had a modest (about 20%) increase in the relative risk of breast cancer compared with women who had never used oral contraceptives. The risk increase varied from 0% to 60%, depending on the specific type of oral combined hormone contraceptive. The risk of breast cancer also increased the longer oral contraceptives were used.

- **Cervical cancer:** Women who have used oral contraceptives for 5 or more years have a higher risk of cervical cancer than women who have never used oral contraceptives. The longer a woman uses oral contraceptives, the greater the increase in her risk of cervical cancer. One study found a 10% increased risk for less than 5 years of use, a 60% increased risk with 5–9 years of use, and a doubling of the risk with 10 or more years of use (9). However, the risk of cervical cancer has been found to decline over time after women stop using oral contraceptives (10–12).
- **Endometrial cancer:** Women who have ever used oral contraceptives have a lower risk of endometrial cancer than women who have never used oral contraceptives. Risk is reduced by at least 30%, with a greater risk reduction the longer oral contraceptives were used (13). The protective effect persists for many years after a woman stops using oral contraceptives (12, 14, 15). An analysis of women participating in the prospective NIH-AARP Diet and Health Study found that the risk reduction was especially pronounced in those long-time users of oral contraceptives who were smokers, obese, or exercised rarely (13).
- **Ovarian cancer:** Women who have ever used oral contraceptives have a 30% to 50% lower risk of ovarian cancer than women who have never used oral contraceptives (16–18). This protection has been found to increase with the length of time oral contraceptives are used (13) and to continue for up to 30 years after a woman stops using oral contraceptives (17). A reduction in ovarian cancer risk with use of oral contraceptives is also seen among women who carry a harmful mutation in the *BRCA1* or *BRCA2* gene (19–21).
- **Colorectal cancer:** Oral contraceptive use is associated with 15% to 20% lower risks of colorectal cancer (12, 14, 22, 23).

How could oral contraceptives influence cancer risk?

Naturally occurring estrogen and progesterone stimulate the development and growth of some cancers (e.g., cancers that express receptors for these hormones, such as breast cancer). Because birth control pills contain synthetic versions of these female hormones, they could potentially also increase cancer risk.

In addition, oral contraceptives might increase the risk of cervical cancer by changing the susceptibility of cervical cells to persistent infection with high-risk HPV types (the cause of virtually all cervical cancers).

Researchers have proposed multiple ways that oral contraceptives may lower the risks of some cancers, including:

- suppressing endometrial cell proliferation (endometrial cancer)
- reducing the number of ovulations a woman experiences in her lifetime, thereby reducing exposure to naturally occurring female hormones (ovarian cancer)
- lowering the levels of bile acids in the blood for women taking oral conjugated estrogens (colorectal cancer) (23)

Selected References

1. Burkman R, Schlesselman JJ, Ziemann M. Safety concerns and health benefits associated with oral contraception. *American Journal of Obstetrics and Gynecology* 2004; 190(4 Suppl):S5–22. [PubMed Abstract]
2. Bassuk SS, Manson JE. Oral contraceptives and menopausal hormone therapy: relative and attributable risks of cardiovascular disease, cancer, and other health outcomes. *Annals of Epidemiology* 2015; 25(3):193–200 [PubMed Abstract]
3. Wentzensen N, Berrington de Gonzalez A. The Pill's gestation: from birth control to cancer prevention. *Lancet Oncology* 2015; 16(9):1004–6. doi: 10.1016/S1470-2045(15)00211-9
4. Collaborative Group on Hormonal Factors in Breast Cancer. Breast cancer and hormonal contraceptives: collaborative reanalysis of individual data on 53,297 women with breast cancer and 100,239 women without breast cancer from 54 epidemiological studies. *Lancet* 1996; 347(9017):1713–1727. [PubMed Abstract]
5. Hunter DJ, Colditz GA, Hankinson SE, et al. Oral contraceptive use and breast cancer: a prospective study of young women. *Cancer Epidemiology Biomarkers and Prevention* 2010; 19(10):2496–2502. [PubMed Abstract]
6. Bhupathiraju SN, Grodstein F, Stampfer MJ, et al. Exogenous hormone use: Oral contraceptives, postmenopausal hormone therapy, and health outcomes in the Nurses' Health Study. *American Journal of Public Health* 2016; 106(9):1631–1637. [PubMed Abstract]

7. Beaber EF, Buist DS, Barlow WE, et al. Recent oral contraceptive use by formulation and breast cancer risk among women 20 to 49 years of age. *Cancer Research* 2014; 74(15):4078-4089. [[PubMed Abstract](#)]
8. Mørch LS, Skovlund CW, Hannaford PC, et al. Contemporary hormonal contraception and the risk of breast cancer. *New England Journal of Medicine* 2017; 377(23):2228-2239. [[PubMed Abstract](#)]
9. Smith JS, Green J, Berrington de Gonzalez A, et al. Cervical cancer and use of hormonal contraceptives: a systematic review. *Lancet* 2003; 361(9364):1159-1167. [[PubMed Abstract](#)]
10. International Collaboration of Epidemiological Studies of Cervical Cancer, Appleby P, Beral V, et al. Cervical cancer and hormonal contraceptives: collaborative reanalysis of individual data for 16,573 women with cervical cancer and 35,509 women without cervical cancer from 24 epidemiological studies. *Lancet* 2007; 370(9599):1609-1621. [[PubMed Abstract](#)]
11. Roura E, Travier N, Waterboer T, et al. The influence of hormonal factors on the risk of developing cervical cancer and pre-cancer: Results from the EPIC Cohort. *PLoS One* 2016; 11(1):e0147029. [[PubMed Abstract](#)]
12. Iversen L, Sivasubramaniam S, Lee AJ, Fielding S, Hannaford PC. Lifetime cancer risk and combined oral contraceptives: The Royal College of General Practitioners' Oral Contraception Study. *American Journal of Obstetrics and Gynecology* 2017; 216(6):580.e1-580.e9. [[PubMed Abstract](#)]
13. Michels KA, Pfeiffer RM, Brinton LA, Trabert B. Modification of the associations between duration of oral contraceptive use and ovarian, endometrial, breast, and colorectal cancers. *JAMA Oncology* 2018; doi:10.1001/jamaoncol.2017.4942. [[PubMed Abstract](#)]
14. Gierisch JM, Coeytaux RR, Urrutia RP, et al. Oral contraceptive use and risk of breast, cervical, colorectal, and endometrial cancers: a systematic review. *Cancer Epidemiology, Biomarkers & Prevention* 2013; 22(11):1931-1943. [[PubMed Abstract](#)]
15. Collaborative Group on Epidemiological Studies on Endometrial Cancer. Endometrial cancer and oral contraceptives: an individual participant meta-analysis of 27 276 women with endometrial cancer from 36 epidemiological studies. *Lancet Oncology* 2015; 16(9):1061-1070. [[PubMed Abstract](#)]
16. IARC Working Group on the Evaluation of Carcinogenic Risks to Humans. Pharmaceuticals. [Combined estrogen-progestogen contraceptives](#). *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans* 2012; 100A:283-311.
17. Havrilesky LJ, Moorman PG, Lowery WJ, et al. Oral contraceptive pills as primary prevention for ovarian cancer: a systematic review and meta-analysis. *Obstetrics and Gynecology* 2013; 122(1):139-147. [[PubMed Abstract](#)]
18. Wentzensen N, Poole EM, Trabert B, et al. Ovarian cancer risk factors by histologic subtype: An analysis from the Ovarian Cancer Cohort Consortium. *Journal of Clinical Oncology* 2016; 34(24):2888-2898. [[PubMed Abstract](#)]

19. Iodice S, Barile M, Rotmensz N, et al. Oral contraceptive use and breast or ovarian cancer risk in BRCA1/2 carriers: a meta-analysis. *European Journal of Cancer* 2010; 46(12):2275-2284. [\[PubMed Abstract\]](#)
20. Moorman PG, Havrilesky LJ, Gierisch JM, et al. Oral contraceptives and risk of ovarian cancer and breast cancer among high-risk women: a systematic review and meta-analysis. *Journal of Clinical Oncology* 2013; 31(33):4188-4198. [\[PubMed Abstract\]](#)
21. Friebel TM, Domchek SM, Rebbeck TR. Modifiers of cancer risk in BRCA1 and BRCA2 mutation carriers: systematic review and meta-analysis. *Journal of the National Cancer Institute* 2014; 106(6):dju091. [\[PubMed Abstract\]](#)
22. Luan NN, Wu L, Gong TT, et al. Nonlinear reduction in risk for colorectal cancer by oral contraceptive use: a meta-analysis of epidemiological studies. *Cancer Causes & Control* 2015; 26(1):65-78. [\[PubMed Abstract\]](#)
23. Murphy N, Xu L, Zervoudakis A, et al. Reproductive and menstrual factors and colorectal cancer incidence in the Women's Health Initiative Observational Study. *British Journal of Cancer* 2017; 116(1):117-125. [\[PubMed Abstract\]](#)

Related Resources

[Breast Cancer Prevention \(PDQ®\)—Patient Version](#)

[Cervical Cancer Prevention \(PDQ®\)—Patient Version](#)

[Colorectal Cancer Prevention \(PDQ®\)—Patient Version](#)

[Endometrial Cancer Prevention \(PDQ®\)—Patient Version](#)

[HPV and Cancer](#)

[Ovarian, Fallopian Tube, and Primary Peritoneal Cancer Prevention \(PDQ®\)—Patient Version](#)

Reviewed: February 22, 2018

If you would like to reproduce some or all of this content, see [Reuse of NCI Information](#) for guidance about copyright and permissions. In the case of permitted digital reproduction, please credit the National Cancer Institute as the source and link to the original NCI product using the original product's title; e.g., "Oral Contraceptives and Cancer Risk was originally published by the National Cancer Institute."